

Patients COVID-19 Vaccine After-Care Sheet

Pfizer/Moderna vaccine

For providers			回旅
Date:			7 .70
Today I received the (circle one):	Pfizer	Moderna	(a) (b)
Today I received dose number:			
Future dose appointment details (if applicable):			Scan to lead



-19 vaccines

It's great you got your vaccination today!

Vaccination will help protect you from serious COVID-19 illness. After vaccination, many people will have some symptoms which are a sign that your body is working with the vaccine. These symptoms may make you feel like you have the flu and may make it difficult to do some of your regular activities for 24-48 hours.

Most people will have one or a few of these side effects

Pain at injection site Muscle pain or joint pain Tiredness Upset stomach, vomiting or

diarrhea Headache Chills Fever

Some people may have one or a few of these side effects

Loss of appetite Swollen lymph nodes Dizziness or sleepiness Rash or itchy skin (not at injection site) **Excessive** sweating

If you have any of these serious symptoms, call 9-1-1

Swelling of your face or mouth Hives

Trouble breathing Very pale colour and

serious drowsiness

High fever (over 40°c)

Convulsions or seizures Other serious symptoms like "pins and needles" or numbness

How to treat side effects

- To reduce pain or discomfort at the injection site, apply a clean, cool, wet washcloth over the area and lightly move your arm.
- For muscle pain and fever, you can take over-the-counter medications like Tylenol and Advil, unless there are other reasons you shouldn't take those. If you're not sure, ask your primary care provider. If you start with Tylenol or Advil and the fever doesn't go down to below 37.8°c after an hour, it's safe to try the other one an hour later to see if it works better.
- If you have fever or diarrhea, make sure to drink plenty of fluids.
- If you have fever, rest and dress in light clothing to keep you cool.







If side effects aren't going away after 48 hours, call: Telehealth: 1-866-797-0000 or your Primary Care Provider:

Remember

- · You will need a full series of shots, as recommended by your local public health authorities, to be optimally protected. Even if you have side effects from the first shot, it's important to get the remaining shots unless your primary care provider or vaccination provider tells you not to.
- · Your first dose helps to protect you from serious cases of COVID-19, but you will not be optimally protected until 2 to 4 weeks after your last dose.
- Some groups of people are eligible for a third dose to help protect them against COVID-19. Please speak to your primary care provider or vaccination provider for more information.
- · You MUST continue to follow public health measures. Even if you're vaccinated, you could still pass the virus on to someone who isn't vaccinated yet. Cover your mouth and nose with a mask when around others, stay at least 6 feet away from others, avoid crowds, and wash your hands often.
- · You can help the vaccine effort! Tell your friends and family you're happy to have gotten the vaccine and that you hope that they will get their vaccine soon too.



COVID-19 Vaccine After-Care Sheet is a product of the Centre for Effective Practice to use, copy, and distribute this material for all non-commercial and research purposes is granted, provided this paragraph and appropriate citations appear in all copies, modifications, and distributions. COVID-19 Vaccine After-Care Sheet for commercial purposes or any modifications of the Resource are subject to charge and use must be negotiated with the Centre for Effective Practice (Email: info@cep.health).

